HUMATES

NTS
Nutri-Tech Solutions®

FAST FULVIC™

High analysis liquid fulvic acid

An 8% fulvic acid liquid possessing a myriad of yield-enhancing qualities.

Fulvic Acid is a powerful organic extract from humates which offers a host of benefits for both the soil and the plant. This is a multi-functional chelating agent, microbe promotant and growth enhancer all in one.

KEY PERFORMANCE

- High analysis liquid fulvic acid
- Powerful natural chelating agent
- Stabilise fertilisers & magnify response
- Promote beneficial soil microbes
- Increases nutrient uptake
- Improves availability of rock minerals
- Increases crop yield and quality

GROWTH STAGE

- ✓ Pre-planting
- ✓ Planting
- ✓ Vegetative Growth
- ✓ Pre-flower, Flowering, Fruit Set & Fill

SIZES:

1L, 5 L, 20 L, 200 L, 1000 L

PRODUCT CODE:

NTSFF

ACO Certification No. 456

TYPICAL ANALYSIS W/V

Fulvic Acid	8%
SG	1.1
pH	3.5 – 4.0
Conductivity	20 – 40 mS/cm
Appearance	Brown liquid



FAST FULVIC™

APPLICATION RATES

Horticulture & Turf

FOLIAR SPRAY RATE
0.5 L per 100 L water up to 5 L/ha equivalent

FERTIGATION RATE
3 – 5 L/Ha

FREQUENCY
Weekly or as required. Include with all nutrient foliar sprays (confirm compatibility)

Broadacre Crops

FOLIAR SPRAY RATE
1 L/ha in 60 – 100 L water, depending on canopy closure

FREQUENCY
Include with all nutrient foliar sprays (confirm compatibility)

Pasture

FOLIAR SPRAY RATE
1 – 2 L/ha in 60 – 100 L water

FREQUENCY
Include with all nutrient foliar sprays (confirm compatibility)

Spot-Spraying

FOLIAR SPRAY RATE
50 mL/ha per 10 L water

FREQUENCY
As required

*Disclaimer: Rates are dependent on biomass and water rates and should be used as a guide only. Application rates and frequencies should be adjusted according to crop monitoring and demand. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare as listed on label. Sensitive plants, or foliage/fruit easily susceptible to staining, should be test sprayed prior to full scale application. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application. Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use. Refer to product label or PIS for alternate application rates.

Nutri-Tech Solutions P/L (NTS)
7 Harvest Road, Yandina, Queensland 4561, Australia
P +61 7 5472 9900 | **E** info@nutri-tech.com.au
www.nutri-tech.com.au

INSTRUCTIONS

Stir well before use.

Rates and frequencies should be adjusted according to crop monitoring.

Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare. Sensitive foliage or plants should be test sprayed prior to full scale application.

Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application.

IMPORTANT

Compatibility and/or performance cannot be guaranteed when combining with other products (including various herbicide formulations).

Jar test for compatibility and spray on a small area before widespread application. Avoid spraying close to harvest if produce is susceptible to staining.

Note: Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.

SAFETY DIRECTIONS

May cause skin irritation. Keep locked up and out of reach of children. Avoid contact with skin. Wear suitable gloves.

Additional information is listed in the Safety Data Sheet.

STORAGE & HANDLING

Store in a cool dry area out of direct sunlight.

Do not pre-mix or store in diluted form.

Seal lid/cap immediately after use.

YOUR LOCAL NTS DISTRIBUTOR:

biologix

201 New Renwick Road, Burleigh
Blenheim 7272, New Zealand
P +64 21 190 1051
E info@biologix.co.nz
W biologix.co.nz

DEC 2023